Microbial source tracking An overview and potential advances with microarray technology

Douglas R. Call
Dept. Veterinary Microbiology & Pathology
Washington State University
Pullman, WA
drcall@wsu.edu

Why source tracking?

- Management and mitigation
- Better risk assessment

Possible markers

- Bacteria E. coli, Enterococcus,
 Bacteroides, Bifidobacterium
- Protozoa Cryptosporidium spp. oocysts
- Virus coliphages
- Chemical (caffeine, fecal steroids)
- Immunological targets (sIgA)

Microbial Source Tracking (MST)

- Assumptions:
 - Host specificity exists
 - Ratios remain constant
 - No significant environmental replication
 - No significant environmental reservoirs

Strategies with bacteria

Library dependent


Antimicrobial resistance


Phenotype PFE g., antibiotic Ribotyping, RAPDs, DGGE,


AFLPs


Genotype, e.g.,
Library independent esence differention of specific genes or "fingerprints"

Microarrays










Wiggins et al. 2003. Use of antibiotic resistance analysis for representativeness testing of multiwatershed libraries. (Applied and Environmental Microbiology, 2003)

- Six watersheds (Virginia)
- 6,587 Enterococci
- Human, domesticated and wild animal sources
- Characterized using 10 drugs (3 levels)

Source (n=6,587)	Human	Domestic	Wild
Human (n=1,970)	63%	19%	18%
Domestic (n=3,345)	20%	54%	26%
Wild (1,272)	21%	24%	55%




<u>Challenges</u>


- Technical expertise
- Practical application
 - Sampling regime
 - Assay cost
 - Sensitivity, specificity
 - Assay format
 - Validation

Strategies with bacteria


- Library dependent
 - Antimicrobial resistance analysis
 - REP-PCR, PFGE,
 Ribotyping, RAPDs, DGGE
 AFLPs
- Library independent
 - Host-specific genetic markers
- Microarrays








What is a microarray?




Probes are "printed" on the slide


Targets are detected after hybridization













Mixed-Genome Microarrays



Example hybridization with mixed-genome, Enterococcus microarray





(n = 4,320 probes)


Human isolate

Elk/deer isolate

Cow isolate








- Data normalized by subgrid using 16S rDNA probe
- Preliminary data includes 83 hybridizations
- N = 358,560 data points


Proportion of putative markers from host-specific libraries



Discriminant function analysis

	Cow	Dog	Elk	Human	Wfowl
Cow	14	1		2	1
Dog	2	12			3
Elk			13		
Human	1			15	1
Wfowl	3	1		1	13

19 probes; 98.5% correct classification rate 8 probes; 75.9% correct classification rate



What's next?

- Work plan:
 - Complete additional 200-250 hybridizations
 - Narrow selection (<400 probes)
 - Validate markers against independent isolates
 - Verify host specificity
 - Quantify marker carriage
 - Test with water samples
 - Develop PCR assays (suspension arrays?)

Summary

Phenotypic or genotypic markers

- Library-dependent
 - Large effort to prepare
 - Validity over space and time?
 - 50-80% correct classification good enough?

Summary

- Library-independent
 - Few markers available
 - Validity still needs to be confirmed
 - Still needs to be cheap and simple

<u>Acknowledgements</u>

Co-investigators:

- •Frank Loge, Civil & Environ. Eng., Pullman
- John Gay, VCS & FDIU, Pullman

Collaborators:

- Dale Hancock, VCS & FDIU, Pullman
- Tom Besser, VMP & WADDL, Pullman
- •Monica Borucki, USDA-ARS, Pullman

Funding:

- •USDA-CSREES (2002-01078) (8/02-7/05)
- Agricultural Animal Health Program, WSU

Staff & others:

- Marilyn Soule, Ph.D.
- Melissa Krug
- Stacey LaFrentz
- Melissa Oatley
- Field Disease
 Investigation Unit

Needed: fecal samples from around the country. Please contact: Doug Call, drcall@wsu.edu, 509-335-6313